

Open Access Article

## Phenolic Content, Antioxidant, Cytotoxic of Fractions of *Spatholobus Littoralis* Hassk from Kalimantan, Indonesia

Dodi Iskandar<sup>1,2</sup>, Nashi Widodo<sup>3</sup>, Warsito<sup>4</sup>, Masruri<sup>4\*</sup>, Rollando<sup>5</sup>, Yeyasa Poulorossi P. Antang<sup>6</sup>

<sup>1</sup> Agricultural Technology Department, Polytechnic State Pontianak, Pontianak 78124, Indonesia

<sup>2</sup> Doctoral Student, Chemistry Department, Brawijaya University, Jl. Veteran, Malang 65145, Indonesia

<sup>3</sup> Biology Department, Brawijaya University, Jl. Veteran, Malang 65145, Indonesia

<sup>4</sup> Chemistry Department, Brawijaya University, Jl. Veteran, Malang 65145, Indonesia

<sup>5</sup> Pharmacy Department, Ma Chung University, Jl. Villa Puncak Tidar Blok N No.1, Malang 65151, Indonesia

<sup>6</sup> Sanggau District Government, West Kalimantan, Jl. RE Martadinata No 14-15, Sanggau 78512, Indonesia

**Abstract:** *Spatholobus littoralis* Hassk (SLH) plant grows wild in the forests of the island of Kalimantan. This plant has the potential as an antioxidant and anticancer. The current research was to quantify the total phenol content, measure the antioxidant and anticancer power against breast cancer cells in non-polar (hexane), polar (water), and semi-polar (ethyl acetate) fractions from *Spatholobus littoralis* wood. Phenol content was determined with gallic acid as standard. Antioxidant activity was measured by the 2,2-diphenyl-1-picrylhydrazyl (DPPH) method, the IC<sub>50</sub> value was determined. Anticancer strength against breast cancer cells was determined using 4T1 cell culture. Toxicity to normal cells was analyzed using Vero cells. The selectivity index (SI) was calculated by dividing the IC<sub>50</sub> of Vero cells by the IC<sub>50</sub> of 4T1 cells. The results showed that total phenol for ethanol extract and water fraction were 350 mg and 146.9 mg equivalent to gallic acid mg/50 g SLH. The IC<sub>50</sub> values of antioxidant were 198.76 ppm, 349.89 ppm, and 2.17 ppm for the hexane, water, and ethyl acetate fractions respectively. Breast anticancer assay with cell line 4T1 showed that the IC<sub>50</sub> values of the hexane fraction and fraction of ethyl acetate in sequence were 20.0 mcg/mL and 7.4 mcg/mL. Toxicity to Vero cells (cell no cancer) showed the IC<sub>50</sub> value for the hexane and ethyl acetate fractions were 131.52 mcg/mL and 79.55 mcg/mL respectively. SI for hexane fractions and ethyl acetate fractions were 6.57 and 10.75 respectively. In summary, the SLH plant has the potential as an antioxidant and anticancer of the breast and can continue in further research. The novelty of this study is the quantification of phenol content, the strength of antioxidant activity, and breast cancer in vitro.

**Keywords:** *spatholobus littoralis* hassk, phenol content, breast anticancer, selectivity index.

### 来自印度尼西亚加里曼丹的石鳖组分的酚类含量、抗氧化剂和细胞毒性

**摘要:** 石鳖植物生长在加里曼丹岛的森林中。这种植物具有抗氧化和抗癌的潜力。目前的研究是量化总酚含量, 测量来自山楂木的非极性(己烷)、极性(水)和半极性(乙酸乙酯)级分中对乳腺癌细胞的抗氧化和抗癌能力。以没食子酸为标准测定苯酚含量。采用2,2-二苯基-1-苦基肼法测定抗氧化活性, 测定半数最大抑菌浓度值。使用4T1细胞培养物确定针对乳腺癌细胞的抗癌强度。使用维罗细胞分析对正常细胞的毒性。通过将维罗细胞的半数最大抑菌浓度除以4T1细胞的半数最大抑菌浓度计算选择性指数。结果表明, 乙醇提取物和水馏

Received: 15 December, 2022 / Revised: 15 January, 2022 / Accepted: 16 February, 2022 / Published: 28 March, 2022

Fund Project: WCR Grant-DIKTI-Indonesia

About the authors: Dodi Iskandar, Agricultural Technology Department, Polytechnic State Pontianak, Pontianak, Indonesia; Doctoral Student, Chemistry Department, Brawijaya University, Malang, Indonesia; Nashi Widodo, Biology Department, Brawijaya University, Malang, Indonesia; Warsito, Chemistry Department, Brawijaya University, Malang, Indonesia; Masruri, Chemistry Department, Brawijaya University, Malang, Indonesia; Rollando, Pharmacy Department, Ma Chung University, Malang, Indonesia; Yeyasa Poulorossi P. Antang, Sanggau District Government, Sanggau, Indonesia

Corresponding author Masruri, iskandar.dodi79@gmail.com

分的总酚分别为350毫克和146.9毫克，相当于没食子酸毫克/50公克山楂。对于己烷、水和乙酸乙酯馏分，抗氧化剂的半数最大抑菌浓度值分别为198.76百万分之几、349.89百万分之几和2.17百万分之几。4T1细胞系乳腺癌抗癌试验表明，己烷组分和乙酸乙酯组分的半数最大抑菌浓度依次为20.0微克/毫升和7.4微克/毫升。对维罗细胞（无癌细胞）的毒性显示，己烷和乙酸乙酯部分的半数最大抑菌浓度值分别为131.52微克/毫升和79.55微克/毫升。己烷馏分和乙酸乙酯馏分的SI分别为6.57和10.75。总之，山楂植物具有作为抗氧化剂和抗乳腺癌的潜力，可以继续进一步的研究。这项研究的新颖之处在于对苯酚含量、抗氧化活性强度和体外乳腺癌的量化。

**关键词：**山楂、酚含量、乳腺癌抗癌、选择性指数。

## 1. Introduction

Plants have been used as medicine by humans for thousands of years. People still use medicinal plants as a source of antioxidants, anticancer, antiviral, antibacterial, and antidote to other diseases [1]. In developing countries, 80 percent of people still depend on medicinal plants to treat diseases and health problems [2]. People are increasingly aware of maintaining health by consuming ingredients from nature that contain antioxidant compounds, antivirals, immune boosters, anticancer, and other active compounds. This awareness has increased sharply, especially during the COVID-19 coronavirus pandemic [3]. The number of herbal purchases during the corona pandemic in Indonesia has increased. This is evidenced by a study conducted in the province of South Kalimantan, Indonesia. The results stated that during the pandemic, 88% of the people consumed products from medicinal plants, namely jamu gendong. Likewise, the production of jamu gendong has increased by 100% [4]. Likewise, in general, in the whole of Indonesia, public consumption of herbal medicines increased by 41% [5]. The popularity of herbal medicines can be boosted by people's experience after consuming and feeling the benefits, besides that it is also supported by scientific data and clinical trials [6]. Although herbal medicines are widely used, the absence of information about the toxicity of herbal medicine can lead to misuse. Thus, the presence of toxicity data will make users feel safe [7].

One of the plants that can be used as traditional medicine is the genus *spatholobus*. Plants of the genus *spatholobus* have many bioactivities. Several publications mention that this genus of plants has efficacy as a hypoallergenic [8], antimicrobial [9, 10], antituberculosis [11], radical scavenging [12], antioxidant [13], cytotoxic [14], antimicrobial [10, 15] and anticancer [16]. Antioxidants are substances in small amounts that can delay or block the oxidation of the substrate [17]. Antioxidants play an important role

in the prevention and treatment of cancer. Cancer therapy using drugs can cure. However, the drug's effectiveness is limited to a few deleterious side effects. Therefore, to erode the negative effects of a drug, the consumption of antioxidants is often necessary. In addition, antioxidants can also reduce the active substance and minimize persistent oxidative damage [18]. However, the exact number of doses given to patients with cancer must be taken seriously [19]. Medicinal plants with high phenolic content are thought to provide antioxidant activity and make a significant contribution to the fight against pathological conditions such as cancer, diabetes, aging, cardiovascular, and other degenerative diseases [20].

Breast cancer is a disease caused by tumors that are the most common in women and the most malignant, and the highest cause of death in more than 100 countries in the world [21]. Based on Globocan 2020, breast cancer patients in Indonesia reached 65,858 new cases and 22,430 deaths [22]. The process that causes failure in treatment due to breast cancer and the death of cancer sufferers is metastatic. Metastatic breast cancer is influenced by two factors, which include intrinsic and extrinsic. Intrinsic are MYC, PI3KA, and TP53 mutations. Extrinsic factors include hypoxia, oxidative stress, and acidosis [23]. To determine the efficacy of a plant drug against metastatic breast cancer, it can be done by in vitro assay using cell lines such as 4T1 [24]. Cell line 4T1 is one of the cell models that is widely used as a representative of breast cancer cells [25].

One species of the genus *Spatholobus* in Indonesia is *Spatholobu littoralis* Hassk (SLH). This plant grows wild in the forests of the island of Kalimantan. Previous research stated that the ethanol extract of this plant has antioxidant activity with an IC<sub>50</sub> value of 8.25 g/ml using the DPPH method [26]. In addition, anticancer test of water extract of *spatholobus littoralis* hassk wood using several cell cultures showed IC<sub>50</sub> 1,063.28 g/mL (MCF7), 53.34 g/mL (HepG2), 150.63

g/mL (T47D), 114.38 µg/mL (WiDR), 97.50 µg/mL (HTB), 182.95 µg/mL (HeLa), and 710.10 µg/mL (Vero) [27]. In this study, we have carried out total phenol assays, antioxidant, anticancer, and toxicity tests in vitro in three groups of non-polar, semi-polar, and polar solvents.

## 2. Materials and Methods

### 2.1. Sample Collection

SLH plants were obtained from the Lamandau district, Central Kalimantan Province, Indonesia and identified in the taxonomy laboratory of the Department of Biology at the University of Tanjungpura on March 3, 2021. The sample used is the wood part. The wood is chopped and made into powder at the Materia Medica laboratory in Batu, Indonesia.

### 2.2. Chemicals

Materials with analytical grades used include distilled water, methanol, dragendorff, mayer, wagner, salkowski, froath, ferric chloride, gelatin alkaline, lead acetate, modified bortrager, molisch, benedict, fehling a&b, ninhydrin, ciulei, gallic acid, Folin-Ciocalteu Reagent, Sodium carbonate, diphenyl picrylhydrazin (DPPH) and ethanol. Technical solutions include ethanol, ethyl acetate, and hexane.

### 2.3. Preparation of SLH Extracts

*Spatholobus littoralis* Hassk Plant identification was carried out in the Biology laboratory of Tanjung Pura University (0.0606° S and 109.344056° E, West Borneo, Indonesia) on March 3, 2021. 2.7 kg SLH wood powder was macerated with technical ethanol (96%), concentrated with a rotary evaporator, and dried in a water bath at 50 Celsius. Obtained dry crude extract as much as 97 grams. The dry extract was crushed in a mortar, dissolved in distilled water, and sonicated into a suspension. Then in the first step, 30 grams of the dry extract was fractionated using the liquid-liquid partition method using water to hexane ratio of 3:1. Then in the second stage, the water fraction was partitioned with ethyl acetate with a ratio of 3:1. In the third stage, the water fraction is obtained. Each fraction was concentrated and dried and obtained 3.7 g, 4.9 g, and 7 g of hexane, ethyl acetate, and water respectively.

### 2.4. Analysis

#### 2.4.1. Qualitative Phytochemical Screening Tests

The method used was adapted with slight modifications [28]. These qualitative tests include alkaloids, phytosterol, saponins, phenols, tannins, flavonoids, glycosides, carbohydrates, protein & amino acids, and triterpenoids.

#### 2.4.2. Total Phenol Content Test

The method used was adapted with slight modifications [29].

##### 2.4.2.1. Standard Calibration Curve

Weighed 50 mg of gallic acid, added 1 mL of 96% ethanol, then added distilled water until the final volume was 50 mL so that a concentration of 1 mg/mL was obtained. From the main solution of gallic acid with a concentration of 1 mg/mL pipette 1 mL, 2 mL, 3 mL, 4 mL, and 5 mL respectively and then diluted with distilled water to a final volume of 10 mL to produce concentrations of 100, 200, 300, 400 and 500 ppm. gallic acid, respectively. From each concentration of the gallic acid solution, 0.1 mL was pipetted and then 7.9 mL of distilled water and 0.5 mL of Folin-Ciocalteu reagent were added and shaken until homogeneous and allowed to stand for 8 minutes. 1.5 mL of 10% Na<sub>2</sub>CO<sub>3</sub> solution was added then shaken homogeneously and then allowed to stand for 2 hours at room temperature. Measure the absorption at a maximum absorption wavelength of 765 nm, then a calibration curve was made for the relationship between gallic acid concentration (µg/ml) and absorption.

##### 2.4.2.2. Total Phenol Content

Each ethanol extract and water extract weighed 100 mg and dissolved to 10 mL with distilled water to obtain a concentration of 10 mg/mL. 1 mL was taken and diluted with distilled water to 10 mL to obtain an extract concentration of 1 mg/mL. Dipiper 0.2 mL of extract and add 15.8 mL of distilled water and 1 mL of Folin-Ciocalteu reagent, then shaken. Let stand for 8 minutes and then add 3 mL of 10% NaCO<sub>3</sub> to the mixture. Allow the mixture to stand for 2 hours at room temperature. Absorption was measured using a UV-Vis spectrophotometer at a wavelength of 765 nm for 3 replications to obtain the phenol content as mg gallic acid/mg material.

#### 2.4.3. In Vitro Antioxidant Activity Test

The method used was adapted with slight modifications [30].

##### 2.4.3.1. DPPH Preparation

Weighed 2 mg of DPPH, dissolved in 50 ml of ethanol, incubated for 30 minutes in a dark room at room temperature, and measured at a wavelength of max.

##### 2.4.3.2. Sample Preparation

The hexane, ethyl acetate, and water fractions were weighed 10 mg each., dissolved in 10 ml of ethanol, made with concentrations 62.5 ppm, 125 ppm 250 ppm, 500 ppm, and 1000 ppm. Furthermore, incubated for 30 minutes in the dark at room temperature. Each

fraction was added DPPH solution with a ratio of 1:1, absorbance was measured using a spectrophotometer at a wavelength of max (712 nm) and calculated % inhibition using with this equation:

$$\%Inhibition = \frac{A_{control} - A_{sampil}}{A_{control}} \times 100\% \quad (1)$$

The standard linear curve is created  $y=ax+b$  and calculated  $IC_{50}$  value ( $y=50$ ).

**2.4.5. In Vitro Cytotoxicity Assay**

The method used was adapted with slight modifications [31]. Breast cancer cells (line 4T1) and Vero cells were cultured using RPMI 1640 medium containing 10% fetal bovine serum (FBS), 1% penicillin/streptomycin, and 1% fungal zone. Then, the cells were inoculated in a 5% CO<sub>2</sub> incubator for 24 h (37 C; 95% relative humidity). Next, 8×10<sup>3</sup> cells in 100 L of the compound were isolated in the test solution (31,25-500 g/mL), then incubated for 24 hours. Next, the cells were washed with PBS and added 100 L of MTT reagent/well, incubated for 4 hours at 37 °C. After that, 100 L of 10% SDS solution was added/well and incubated overnight at room temperature in a dark room, then determined. absorbance with a microplate reader at 595 nm. Single treatment data were converted to percent viability and used to calculate IC<sub>50</sub>. Percent cell viability is

calculated using the following equation:

$$\%Cell\ viability = \frac{A_{sampil}}{A_{control}} \times 100\% \quad (2)$$

**2.4.6 Selectivity index (SI)**

SI value is determined based on the following equation [32, 33]:

$$SI = \frac{IC_{50}^{no\ cancer\ cell}}{IC_{50}^{cancer\ cell}} \quad (3)$$

**2.5 Statistical Analysis**

Quantitative experiments used measurements three times and the average value was taken. Linear regression equations and graphs using Microsoft Excel tools.

**3. Results and Discussion**

**3.1 Qualitative Phytochemical Content and Total Phenol Content**

Table 1 shows Qualitative phytochemical screening test results where (-) = absent, (+) = present, SLH = *Spatholobus littoralis* Hassk Bark. Table 2 and Fig. 1 show standard calibration curve.

Table 1 Qualitative phytochemical screening test results

no	Metabolites	test	SLH extract			
			hexane	Ethyl acetate	ethanol	aquades
1	Alkaloids	dragendorff	(-)	(-)	(-)	(-)
		Mayer	(-)	(-)	(-)	(-)
		wagner	(-)	(-)	(-)	(-)
2	phytosterol	salkowski	(-)	(+)	(-)	(-)
3	Saponin	Froath	(-)	(-)	(+)	(+)
4	Phenol	Ferric chloride	(-)	(-)	(+)	(+)
5	Tannin	gelatin	(-)	(-)	(-)	(-)
6	Flavonoid	alkaline	(-)	(-)	(-)	(-)
7	Flavonoid	lead acetat	(-)	(-)	(+)	(+)
8	Glycosides	Modified	(-)	(-)	(-)	(-)
		borntrager	(-)	(-)	(-)	(-)
		Molisch	(-)	(-)	(-)	(-)
		Benedict	(-)	(-)	(-)	(-)
10	Protein & amino acids	Fehling A&B	(-)	(-)	(-)	(-)
		Ninhydrin	(-)	(-)	(+)	(+)
11	triterpenoids	Ciulei	(-)	(-)	(+)	(+)




Fig. 1 Standard calibration curve

Table 2. Standard calibration curve

ppm	Absorbance
100	0.121
200	0.200
300	0.259
400	0.390
500	0.430

The absorbance of the ethanol extract sample was measured three times, namely 0.317; 0.318, and 0.317. So the average absorbance of the ethanol extract sample was 0.317. Measurements of water extract absorbance were 0.133; 0.133 and 0.134. So that the

average absorbance of the aqueous extract. The sample was 0.133.

### 3.2. Antioxidant Activity

Fig. 2 and table 3 elaborate antioxidant activity.

Table 3 Antioxidant test of fractions

Fractions	IC <sub>50</sub> (ppm)
Hexane	198.76
Water	349.89
Ethyl acetate	2.17




Fig. 2 DPPH Standard curve of A (hexane fraction), B (water fraction), and C (fraction of ethyl acetate)

### 3.3. Breast Anti-Cancer Bioactivity




Fig. 3 Breast anti-cancer test of fractions (A: hexane; B: water; C: ethyl acetate) against 4T1 cell line




Fig. 4 Cytotoxic test of fractions (A: hexane; B: water; C: ethyl acetate) against Vero cell line.

Table 4 Breast anticancer bioactivity of fractions

Fractions	IC <sub>50</sub> (mcg/mL)		Selectivity Index
	4T1 Cell line	Vero cell	
Hexane	20.0	131.52	6.57
Water	—	—	—
Ethyl acetate	7.4	79.55	10.75

Fig. 3-4 and table 4 elaborate breast anti-cancer bioactivity. Extraction can be done from solid to liquid, from liquid to liquid, and so on. This process transfers a collection of compounds from one phase to another that is physically separated. This is where further process analysis can be carried out [34]. Ethanol and hexane are miscible and indistinguishable from each other.[35]. For this reason, it is not possible to obtain a non-polar fraction such as hexane if the dry extract is first dissolved in ethanol. Thus, by taking advantage of the water's reluctance to mix with hexane, the dry extract of SLH was first dissolved in water to obtain the non-polar (hexane) fraction easily. Furthermore, after the hexane fraction is separated, the water fraction can be partitioned with ethyl acetate solvent which is semi-polar and easily separated from the water fraction, although it requires a long separation time than the separation of the water fraction and hexane fraction.

SLH wood extract qualitatively contained phytosterols in the fraction of ethyl acetate, saponins in the (ethanol and water) fraction, phenol in the ethanol and water fractions, proteins & amino acids in the ethanol and water fractions, and triterpenoids in the ethanol and water fractions. Based on a qualitative phytochemical test, only the ethanol extract and water fraction contained phenol. By using equation  $y = 0.0008x + 0.0376$  (Fig.1) and inserting the absorbance values, total phenol for ethanol extract and water fraction are 350 mg and 146.9 mg equivalent to gallic acid mg/50 g SLH respectively. many accept the opinion that a high total phenol content does not necessarily have a high antioxidant activity as well. This is because each extract has a different capacity of electron donor [36]. The absence of a relationship between the total phenol content and antioxidants can be understood from several perspectives. First, the total phenolic fraction has not combined all the antioxidant compounds. Second, the synergistic interaction between antioxidant compounds in the mixture does not have to depend on the concentration of the mixture but can depend on the structure and properties of the antioxidants themselves. Third, the phenolic content of plants depends on several genetic, environmental, and processing factors [37].

By observing the DPPH standard curve, the hexane and water fractions were able to detect the absorbance value of the solution with concentrations between the ranges of 63.5 – 1000 ppm. In contrast to the fraction of ethyl acetate, it can only be detected in the range of 1-10 ppm. Entering 50 as y from the regression equation for each curve, the  $IC_{50}$  can be obtained. The  $IC_{50}$  values were 198.76 ppm, 349.89 ppm, and 2.17 ppm for the hexane, water, and ethyl acetate fractions respectively (Fig.2). Based on the  $IC_{50}$  value, the antioxidant power can be grouped into 5 categories. First, the activity is very strong if it is < 10 ppm, the activity is strong if it is between 10-50 ppm, the

activity is moderate if it is between 50-100 ppm, less active if it is between 100-250 ppm, and inactive if it is > 250 ppm [38]. Thus, the hexane fraction has less active antioxidant activity, the water fraction is not active and the fraction of ethyl acetate is very strong. It is different from SLH ethanol extract, its antioxidant activity is very strong with an  $IC_{50}$  of 8.25 ppm [26].

In the breast anticancer test using the 4T1 cell line, the greater the concentration of the fraction, the lower the % viability. Because the more active substances in the fraction, the more 4T1 cancer cells will die. This trend can be seen in the experimental results of the hexane fraction and fraction of ethyl acetate. However, this does not apply to the water fraction. It may be that the anticancer substances in the ethanol extract have all shifted to non-polar (hexane) and semi-polar (ethyl acetate) fractions (Fig 3). Breast anticancer assay with cell line 4T1 showed that the  $IC_{50}$  values of the hexane fraction and fraction of ethyl acetate in sequence were 20.0 mcg/mL and 7.4 mcg/mL. several categories of extract cytotoxicity strength based on its  $IC_{50}$  value. very strong <10 mcg/mL, strong between 10-100 mcg/mL, moderate between 100-500 mcg/mL. There are three categories of extract cytotoxicity strength based on its  $IC_{50}$  value (very strong <10 mcg/mL, strong between 10-100 mcg/mL, and moderate between 100-500 mcg/mL) [39]. From these data (Table 4), the cytotoxicity of the hexane fraction is strong and the fraction of ethyl acetate is very strong. Toxicity to Vero cells (cell no cancer) showed the  $IC_{50}$  value for the hexane and ethyl acetate fractions were 131.52 mcg/mL and 79.55 mcg/mL respectively. The selectivity index is defined as the ratio of the toxic concentration of a sample to its effective bioactive concentration [40]. To evaluate the anticancer activity of a sample, the cytotoxicity to non-malignant cell lines must be calculated [32, 33]. SI for hexane fractions and ethyl acetate fractions were 6.57 and 10.75 respectively. Some researchers stated that a sample with SI value  $\geq 10$  then as the chosen one for further research [32]. Others argue that samples with SI values less than 10 and  $\geq 3$  are considered prospective anticancer [41]. Thus, both fractions deserve to be studied further and are prospective as herbs in terms of breast cancer anticancer perspective.

#### 4. Conclusion

Measurable phenol content in ethanol extract and water fraction from SLH wood was 350 mg and 146.9 mg equivalent to gallic acid mg/50 g SLH. The strongest antioxidant activity was the fraction of ethyl acetate ( $IC_{50}$  2.17 ppm) followed by the hexane fraction ( $IC_{50}$  198.76 ppm) and the water fraction ( $IC_{50}$  349.89 ppm). Meanwhile, the greatest strength of breast anticancer activity in vitro was the fraction of ethyl acetate ( $IC_{50}$  7.4 mcg/mL) followed by the hexane fraction ( $IC_{50}$  20.0 mcg/mL). the toxicity of the

hexane and ethyl acetate fractions were 131.52 mcg/mL and 79.55 mcg/mL. so that the selectivity indices for both hexane fractions and ethyl acetate fractions were 6.57 and 10.75 respectively. Both the ethyl acetate and water fractions have anticancer activity in the breast that deserve to be studied further and in more depth.

## References

- [1] MARRELLI M. Medicinal plants. *Plants*, 2021, 10(7): 1355. <https://doi.org/10.3390/plants10071355>
- [2] OKOYE T. C., UZOR P. F., ONYETO C. A., and OKEREKE E. K. Safe African Medicinal Plants for Clinical Studies. In: *Toxicological Survey of African Medicinal Plants*. Elsevier, Amsterdam, 2014: 535–555. <https://doi.org/10.1016/B978-0-12-800018-2.00018-2>
- [3] GALANAKIS C. M., ALDAWOUD T. M. S., RIZOU M., ROWAN N. J., and IBRAHIM S. A. Food Ingredients and Active Compounds against the Coronavirus Disease (COVID-19) Pandemic: A Comprehensive Review. *Foods*, 2020, 9(11): 1701. <https://doi.org/10.3390/foods9111701>
- [4] LESMAYATI S., QOMARIAH R., AWANIS, and PRAMUDYANI L. The impact of Covid-19 pandemic on people's behavior and herbal drink (jamu) processing businesses in Banjarbaru, South Kalimantan. *E3S Web of Conferences*, 2021, 306: 02046. <https://doi.org/10.1051/e3sconf/202130602046>
- [5] HARFIANI E., PUSPITA R., and RAMADHANI I. Characteristics of Indonesian Society in Utilizing Herbs for Covid Prevention during the Covid-19 Pandemic. *IOP Conference Series: Earth and Environmental Science*, 2021, 913(1): 012104. <https://doi.org/10.1088/1755-1315/913/1/012104>
- [6] JAHROMI B., PIRVULESCU I., CANDIDO K. D., and KNEZEVIC N. N. Herbal Medicine for Pain Management: Efficacy and Drug Interactions. *Pharmaceutics*, 2021, 13(2): 251. <https://doi.org/10.3390/pharmaceutics13020251>
- [7] MENSAH M. L. K., KOMLAGA G., FORKUO A. D., FIREMPONG C., ANNING A. K., and DICKSON R. A. Toxicity and Safety Implications of Herbal Medicines Used in Africa. In: *Herbal Medicine*. IntechOpen, London, 2019.
- [8] MIZUGUCHI H., NARIAI Y., KATO S., NAKANO T., KANAYAMA T., KASHIWADA Y., NEMOTO H., KAWAZOE K., TAKAISHI Y., KITAMURA Y., TAKEDA N., and FUKUI H. Maackiain is a novel antiallergic compound that suppresses transcriptional upregulation of the histamine H1 receptor and interleukin-4 genes. *Pharmacology Research & Perspectives*, 2015, 3(5): e00166. <https://doi.org/10.1002/prp2.166>
- [9] WILLIAMS D., PERRY D., CARRAWAY J., SIMPSON S., UWAMARIYA P., and CHRISTIAN O. E. Antigonococcal Activity of (+)-Medicarpin. *ACS Omega*, 2021, 6(23): 15274–15278. <https://doi.org/10.1021/acsomega.1c01590>
- [10] LI B., ZHANG D., ZHANG Y., JIANG D., LI S., LEI W., WANG H., and LIN F. Synthesis and evaluation of novel benzene-ethanol bearing 1,2,4-triazole derivatives as potential antimicrobial agents. *Medicinal Chemistry Research*, 2017, 26(1): 44–51. <https://doi.org/10.1007/s00044-016-1724-6>
- [11] HASAN N., OSMAN H., MOHAMAD S., CHONG W. K., AWANG K., and ZAHARILUDDIN A. S. M. The Chemical Components of *Sesbania grandiflora* Root and Their Antituberculosis Activity. *Pharmaceutics*, 2012, 5(8): 882–889. <https://doi.org/10.3390/ph5080882>
- [12] NOMAN L., OKE-ALTUNTAS F., ZELLAGUI A., YAGLIOGLU A. S., DEMIRTAS I., CARDOSO S. M., AKKAL S., GHERRAF N., and RHOUATI S. A novel benzimidazole and other constituents with antiproliferative and antioxidant properties from *Thymelaea microphylla* Coss. et Dur. *Natural Product Research*, 2017, 31(17): 2032–2041. <https://doi.org/10.1080/14786419.2016.1274888>
- [13] SILVA A. S., REBOREDO-RODRÍGUEZ P., SÚNTAR I., SUREDA A., BELWAL T., LOIZZO M. R., TUNDIS R., SOBARZO-SANCHEZ E., RASTRELLI L., FORBES-HERNANDEZ T. Y., BATTINO M., FILOSA R., DAGLIA M., NABAVI S. F., and NABAVI S. M. Evaluation of the status quo of polyphenols analysis: Part I — phytochemistry, bioactivity, interactions, and industrial uses. *Comprehensive Reviews in Food Science and Food Safety*, 2020, 19(6): 3191–3218. <https://doi.org/10.1111/1541-4337.12629>
- [14] GUO J. Two New Diterpenoids from the Buds of *Wikstroemia Chamaedaphne*. *Molecules*, 2012, 17(6): 6424–6433. <https://doi.org/10.3390/molecules17066424>
- [15] SON N. T., HARADA K., CUONG N. M., and FUKUYAMA Y. Two New Carboxyethylflavanones from the Heartwood of *Dalbergia tonkinensis* and Their Antimicrobial Activities. *Natural Product Communications*, 2017, 12(11): 1934578X1701201. <https://doi.org/10.1177/1934578X1701201115>
- [16] EMAMI S., & GHANBARIMASIR Z. Recent advances of chroman-4-one derivatives: Synthetic approaches and bioactivities. *European Journal of Medicinal Chemistry*, 2015, 93: 539–563. <https://doi.org/10.1016/j.ejmech.2015.02.048>
- [17] SANTOS-SÁNCHEZ N. F., SALAS-CORONADO R., VILLANUEVA-CAÑONGO C., and HERNÁNDEZ-CARLOS B. Antioxidant Compounds and Their Antioxidant Mechanism. In: *Antioxidants*. IntechOpen, London, 2019.
- [18] FERDOUS U. T., & YUSOF Z. N. B. Medicinal Prospects of Antioxidants From Algal Sources in Cancer Therapy. *Frontiers in Pharmacology*, 2021, 12: 593116. <https://doi.org/10.3389/fphar.2021.593116>
- [19] GRIÑAN-LISON C., BLAYA-CÁNOVAS J. L., LÓPEZ-TEJADA A., ÁVALOS-MORENO M., NAVARRO-OCÓN A., CARA F. E., GONZÁLEZ-GONZÁLEZ A., LORENTE J. A., MARCHAL J. A., and GRANADOS-PRINCIPAL S. Antioxidants for the Treatment of Breast Cancer: Are We There Yet? *Antioxidants*, 2021, 10(2): 205. <https://doi.org/10.3390/antiox10020205>
- [20] YU M., GOUVINHAS I., ROCHA J., and BARROS A. I. R. N. A. Phytochemical and antioxidant analysis of medicinal and food plants towards bioactive food and pharmaceutical resources. *Scientific Reports*, 2021, 11(1): 10041. <https://doi.org/10.1038/s41598-021-89437-4>
- [21] TIRUNEH M., TESFAW A., and TESFA D. Survival and Predictors of Mortality among Breast Cancer Patients in Northwest Ethiopia: A Retrospective Cohort Study. *Cancer Management and Research*, 2021, 13: 9225–9234. <https://doi.org/10.2147/CMAR.S339988>
- [22] THE GLOBAL CANCER OBSERVATORY. *Indonesia*, 2020. <https://gco.iarc.fr/today/data/factsheets/populations/360-indonesia-fact-sheets.pdf>
- [23] WANG L., ZHANG S., and WANG X. The Metabolic



- Mechanisms of Breast Cancer Metastasis. *Frontiers in Oncology*, 2021, 10 :602416. <https://doi.org/10.3389/fonc.2020.602416>
- [24] ABDEL-SALAM M. A. L., PINTO B., CASSALI G., BUENO L., PÊGAS G., OLIVEIRA F., SILVA I., KLEIN A., DE SOUZA-FAGUNDES E. M., DE LIMA M. E., and CARVALHO-TAVARES J. LyeTx I-b Peptide Attenuates Tumor Burden and Metastasis in a Mouse 4T1 Breast Cancer Model. *Antibiotics*, 2021, 10(9): 1136. <https://doi.org/10.3390/antibiotics10091136>
- [25] SCHRÖRS B., BOEGEL S., ALBRECHT C., BUKUR T., BUKUR V., HOLTSTRÄTER C., RITZEL C., MANNINEN K., TADMOR A. D., VORMEHR M., SAHIN U., and LÖWER M. Multi-Omics Characterization of the 4T1 Murine Mammary Gland Tumor Model. *Frontiers in Oncology*, 2020, 10. <https://doi.org/10.3389/fonc.2020.01195>
- [26] ISKANDAR D., & WARSIDAH W. Qualitative Phytochemical Screening and Antioxidant Activity of Ethanol Root Extract of *Spatholobus littoralis* Hassk. *The Journal of Food and Medicinal Plants*, 2020, 1(1): 13–15. <https://doi.org/10.25077/jfmp.1.1.13-15.2020>
- [27] YUNIARTI L., KHARISMA Y., RESPATI T., and TEJASARI M. Halal Critical Point Analysis of *Bajakah* Wood (*Spatholobus littoralis* Hassk) Nano Particle as Anticancer Agent. *Global Medical and Health Communication (GMHC)*, 2021, 9(2): 81–87. <https://doi.org/10.29313/gmhc.v9i2.6997>
- [28] MOHAN E., SURIYA S., SHANMUGAM S., and RAJENDRAN K. Qualitative Phytochemical Screening of Selected Medicinal Plants. *Journal of Drug Delivery and Therapeutics*, 2021, 11(2): 141–144. <https://doi.org/10.22270/jddt.v11i2.4609>
- [29] HERNÁNDEZ ZARATE M. S., ABRAHAM JUÁREZ M. R., CERÓN GARCÍA A., OZUNA LÓPEZ C., GUTIÉRREZ CHÁVEZ A. J., SEGOVIANO GARFIAS J. J. N., and AVILA RAMOS F. Flavonoids, phenolic content, and antioxidant activity of propolis from various areas of Guanajuato, Mexico. *Food Science and Technology*, 2018, 38(2): 210–215. <https://doi.org/10.1590/fst.29916>
- [30] YUSOP S. A. T. W., HAFIZI A. S., SABRI W. M. A. W., and RAZIP M. Asaruddin Antioxidant, Antimicrobial and Cytotoxicity Activities of Propolis from Beladin, Sarawak Stingless Bees *Trigona itama* Extract. *Materials Today: Proceedings*, 2019, 19: 1752–1760. <https://doi.org/10.1016/j.matpr.2019.11.213>
- [31] SEGUN P. A., OGBOLE O. O., ISMAIL F. M. D., NAHAR L., EVANS A. R., AJAIYEGBA E. O., and SARKER S. D. Bioassay-guided isolation and structure elucidation of cytotoxic stilbenes and flavonols from the leaves of *Macaranga barteri*. *Fitoterapia*, 2019, 134: 151–157. <https://doi.org/10.1016/j.fitote.2019.02.019>
- [32] PEÑA-MORÁN O., VILLARREAL M., ÁLVAREZ-BERBER L., MENESES-ACOSTA A., and RODRÍGUEZ-LÓPEZ V. Cytotoxicity, Post-Treatment Recovery, and Selectivity Analysis of Naturally Occurring Podophyllotoxins from *Bursera fagaroides* var. *fagaroides* on Breast Cancer Cell Lines. *Molecules*, 2016, 21(8): 1013. <https://doi.org/10.3390/molecules21081013>
- [33] LÓPEZ-LÁZARO M. How many times should we screen a chemical library to discover an anticancer drug? *Drug Discovery Today*, 2015, 20(2): 167–169. <https://doi.org/10.1016/j.drudis.2014.12.006>
- [34] UDAYA SANKAR K. Extraction Processes. In: *Conventional and Advanced Food Processing Technologies*. John Wiley & Sons Ltd, Chichester, 2014: 129–158.
- [35] JUKIĆ I., POŽAR M., and LOVRINČEVIĆ B. Comparative analysis of ethanol dynamics in aqueous and non-aqueous solutions. *Physical Chemistry Chemical Physics*, 2020, 22(41): 23856–23868. <https://doi.org/10.1039/D0CP03160G>
- [36] DIBACTO R. E. K., TCHUENTE B. R. T., NGUEDJO M. W., TIENCHEU Y. M. T., NYOBE E. C., EDOUN F. L. E., KAMINI M. F. G., DIBANDA R. F., and MEDOUA G. N. Total Polyphenol and Flavonoid Content and Antioxidant Capacity of Some Varieties of Persea Americana Peels Consumed in Cameroon. *The Scientific World Journal*, 2021, 2021: 1–11. <https://doi.org/10.1155/2021/8882594>
- [37] KAYAHAN S., & SALOGLU D. Comparison of Phenolic Compounds and Antioxidant Activities of Raw and Cooked Turkish Artichoke Cultivars. *Frontiers in Sustainable Food Systems*, 2021, 5. <https://doi.org/10.3389/fsufs.2021.761145>
- [38] SYABANA M. A., YULIANA N. D., BATUBARA I., and FARDIAZ D. Antidiabetic activity screening and NMR profile of vegetable and spices commonly consumed in Indonesia. *Food Science and Technology*, 2021, 41(1): 254–264. <https://doi.org/10.1590/fst.14120>
- [39] INDRAYANTO G., PUTRA G. S., and SUHUD F. Validation of in-vitro bioassay methods: Application in herbal drug research. *Profiles of Drug Substances, Excipients, and Related Methodology*, 2021, 46: 273–307. <https://doi.org/10.1016/bs.podrm.2020.07.005>
- [40] PRITCHETT J. C., NAESENS L., and MONTOYA J. Treating HHV-6 Infections. In: *Human Herpesviruses HHV-6A, HHV-6B & HHV-7*. Elsevier, Amsterdam, 2014: 311–331.
- [41] WEERAPREEYAKUL N., NONPUNYA A., BARUSRUX S., THITIMETHAROCH T., and SRIPANIDKULCHAI B. Evaluation of the anticancer potential of six herbs against a hepatoma cell line. *Chinese Medicine*, 2012, 7(1): 15. <https://doi.org/10.1186/1749-8546-7-15>

#### 参考文献:

- [1] MARRELLI M. 药用植物。植物, 2021, 10(7): 1355. <https://doi.org/10.3390/plants10071355>
- [2] OKOYE T. C., UZOR P. F., ONYETO C. A., 和 OKEREKE E. K. 用于临床研究的安全非洲药用植物。在: 非洲药用植物毒理学调查。爱思唯尔, 阿姆斯特丹, 2014: 535–555. <https://doi.org/10.1016/B978-0-12-800018-2.00018-2>
- [3] GALANAKIS C. M., ALDAWOUUD T. M. S., RIZOU M., ROWAN N. J., 和 IBRAHIM S. A. 抗冠状病毒病 (新冠肺炎) 大流行的食品成分和活性化合物: 综合综述。食品, 2020, 9(11): 1701. <https://doi.org/10.3390/foods9111701>
- [4] LESMAYATI S., QOMARIAH R., AWANIS, 和 PRAMUDYANI L. 新冠肺炎大流行对南加里曼丹省班贾尔巴鲁的人们行为和草药饮



- 料(查漠)加工企业的影响。E3S会议网络, 2021, 306: 02046. <https://doi.org/10.1051/e3sconf/202130602046>
- [5] HARFIANI E., PUSPITA R., 和 RAMADHANI I. 印度尼西亚社会在新冠肺炎大流行期间利用草药预防冠状病毒病的特点。物理研究所系列会议: 地球与环境科学, 2021, 913(1): 012104. <https://doi.org/10.1088/1755-1315/913/1/012104>
- [6] JAHROMI B., PIRVULESCU I., CANDIDO K. D., 和 KNEZEVIC N. N. 用于疼痛管理的草药: 功效和药物相互作用。药剂学, 2021, 13(2): 251. <https://doi.org/10.3390/pharmaceutics13020251>
- [7] MENSAH M. L. K., KOMLAGA G., FORKUO A. D., FIREMPONG C., ANNING A. K., 和 DICKSON R. A. 非洲使用的草药的毒性和安全性影响。在: 草药。英泰公开赛, 伦敦, 2019.
- [8] MIZUGUCHI H., NARIAI Y., KATO S., NAKANO T., KANAYAMA T., KASHIWADA Y., NEMOTO H., KAWAZOE K., TAKAISHI Y., KITAMURA Y., TAKEDA N., 和 FUKUI H. 麦肯是一种新型抗过敏化合物, 可抑制组胺H1受体和白细胞介素4基因的转录上调。药理学研究与展望, 2015, 3(5): e00166. <https://doi.org/10.1002/prp2.166>
- [9] WILLIAMS D., PERRY D., CARRAWAY J., SIMPSON S., UWAMARIYA P., 和 CHRISTIAN O. E. (+)-药草平的抗淋球菌活性。美国化学会欧米茄, 2021, 6(23): 15274–15278. <https://doi.org/10.1021/acsomega.1c01590>
- [10] LI B., ZHANG D., ZHANG Y., JIANG D., LI S., LEI W., WANG H., 和 LIN F. 作为潜在抗菌剂的新型苯乙醇的合成和评价 1,2,4-三唑衍生物。药物化学研究, 2017, 26(1): 44–51. <https://doi.org/10.1007/s00044-016-1724-6>
- [11] HASAN N., OSMAN H., MOHAMAD S., CHONG W. K., AWANG K., 和 ZAHARILUDDIN A. S. M. 田菁根的化学成分及其抗结核活性。药品, 2012, 5(8): 882–889. <https://doi.org/10.3390/ph5080882>
- [12] NOMAN L., OKE-ALTUNTAS F., ZELLAGUI A., YAGLIOGLU A. S., DEMIRTAS I., CARDOSO S. M., AKKAL S., GHERRAF N., 和 RHOUATI S. 一种来自小叶百里香科的新型苯并咪唑和其他具有抗增殖和抗氧化特性的成分。等杜尔。天然产物研究, 2017, 31(17): 2032–2041. <https://doi.org/10.1080/14786419.2016.1274888>
- [13] SILVA A. S., REBOREDO-RODRÍGUEZ P., SÜNTAR I., SUREDA A., BELWAL T., LOIZZO M. R., TUNDIS R., SOBARZO-SANCHEZ E., RASTRELLI L., FORBES-HERNANDEZ T. Y., BATTINO M., FILOSA R., DAGLIA M., NABAVI S. F., 和 NABAVI S. M. 多酚分析现状评估: 第一部分——植物化学、生物活性、相互作用和工业用途。食品科学与食品安全综合评价, 2020, 19(6): 3191–3218. <https://doi.org/10.1111/1541-4337.12629>
- [14] GUO J. 来自紫花苜蓿芽的两种新二萜。分子, 2012, 17(6): 6424–6433. <https://doi.org/10.3390/molecules17066424>
- [15] SON N. T., HARADA K., CUONG N. M., 和 FUKUYAMA Y. 来自黄檀心材的两种新羧乙基黄酮及其抗菌活性。天然产物通讯, 2017, 12(11): 1934578X1701201. <https://doi.org/10.1177/1934578X1701201115>
- [16] EMAMI S., 和 GHANBARIMASIR Z. 色满4-1衍生物的最新进展: 合成方法和生物活性。欧洲药物化学杂志, 2015, 93: 539–563. <https://doi.org/10.1016/j.ejmech.2015.02.048>
- [17] SANTOS-SÁNCHEZ N. F., SALAS-CORONADO R., VILLANUEVA-CAÑONGO C., 和 HERNÁNDEZ-CARLOS B. 抗氧化化合物及其抗氧化机制。在: 抗氧化剂。英泰克公开赛, 伦敦, 2019.
- [18] FERDOUS U. T., 和 YUSOF Z. N. B. 来自藻类来源的抗氧化剂在癌症治疗中的药用前景。药理学前沿, 2021, 12: 593116. <https://doi.org/10.3389/fphar.2021.593116>
- [19] GRIÑAN-LISON C., BLAYA-CÁNOVAS J. L., LÓPEZ-TEJADA A., ÁVALOS-MORENO M., NAVARRO-OCÓN A., CARA F. E., GONZÁLEZ-GONZÁLEZ A., LORENTE J. A., MARCHAL J. A., 和 GRANADOS-PRINCIPAL S. 用于治疗乳腺癌的抗氧化剂: 我们到了吗? 抗氧化剂, 2021, 10(2): 205. <https://doi.org/10.3390/antiox10020205>
- [20] YU M., GOUVINHAS I., ROCHA J., 和 BARROS A. I. R. N. A. 药用和食用植物对生物活性食品和药物资源的植物化学和抗氧化分析。科学报告, 2021, 11(1): 10041. <https://doi.org/10.1038/s41598-021-89437-4>
- [21] TIRUNEH M., TESFAW A., 和 TESFA D. 埃塞俄比亚西北部乳腺癌患者的生存率和死亡率预测因素: 一项回顾性队列研究。癌症管理和研究, 2021, 13: 9225–9234. <https://doi.org/10.2147/CMAR.S339988>
- [22] THE GLOBAL CANCER OBSERVATORY. 印度尼西亚, 2020. <https://gco.iarc.fr/today/data/factsheets/populations/360-indonesia-fact-sheets.pdf>
- [23] WANG L., ZHANG S., 和 WANG X. 乳腺癌转移的代谢机制。肿瘤学前沿, 2021, 10: 602416. <https://doi.org/10.3389/fonc.2020.602416>
- [24] ABDEL-SALAM M. A. L., PINTO B., CASSALI G., BUENO L., PÊGAS G., OLIVEIRA F., SILVA I., KLEIN A., DE SOUZA-FAGUNDES E. M., DE LIMA M. E., and

- CARVALHO-TAVARES J. LyeTx I-b 肽可减轻小鼠 4T1 乳腺癌模型中的肿瘤负担和转移。抗生素, 2021, 10(9): 1136. <https://doi.org/10.3390/antibiotics10091136>
- [25] SCHRÖRS B., BOEGEL S., ALBRECHT C., BUKUR T., BUKUR V., HOLTSTRÄTER C., RITZEL C., MANNINEN K., TADMOR A. D., VORMEHR M., SAHIN U., 和 LÖWER M. 4T1 鼠乳腺肿瘤模型的多组学表征。肿瘤学前沿, 2020, 10. <https://doi.org/10.3389/fonc.2020.01195>
- [26] ISKANDAR D., 和 WARSIDAH W. 石竹乙醇根提取物的定性植物化学筛选和抗氧化活性。食品和药用植物杂志, 2020, 1(1): 13–15. <https://doi.org/10.25077/jfmp.1.1.13-15.2020>
- [27] YUNIARTI L., KHARISMA Y., RESPATI T., 和 TEJASARI M. 巴加卡木 (石鳖) 纳米粒子作为抗癌剂的清真临界点分析。全球医疗健康传播, 2021, 9(2): 81–87. <https://doi.org/10.29313/gmhc.v9i2.6997>
- [28] MOHAN E., SURIYA S., SHANMUGAM S., 和 RAJENDRAN K. 选定药用植物的定性植物化学筛选。药物输送与治疗学杂志, 2021, 11(2): 141–144. <https://doi.org/10.22270/jddt.v11i2.4609>
- [29] HERNÁNDEZ ZARATE M. S., ABRAHAM JUÁREZ M. R., CERÓN GARCÍA A., OZUNA LÓPEZ C., GUTIÉRREZ CHÁVEZ A. J., SEGOVIANO GARFIAS J. J. N., 和 AVILA RAMOS F. 来自墨西哥瓜纳华托不同地区的蜂胶的黄酮类化合物、酚类含量和抗氧化活性。食品科技, 2018, 38(2): 210–215. <https://doi.org/10.1590/fst.29916>
- [30] YUSOP S. A. T. W., HAFIZI A. S., SABRI W. M. A. W., 和 RAZIP M. 来自贝拉丁的蜂胶的阿萨鲁丁抗氧化、抗菌和细胞毒性活性, 砂拉越无刺蜜蜂螳螂提取物。今日材料: 诉讼, 2019, 19: 1752–1760. <https://doi.org/10.1016/j.matpr.2019.11.213>
- [31] SEGUN P. A., OGBOLE O. O., ISMAIL F. M. D., NAHAR L., EVANS A. R., AJAIYEGBA E. O., 和 SARKER S. D. 生物测定指导的通心粉叶子中细胞毒性二苯乙烯和黄酮醇的分离和结构鉴定。鱼翅目, 2019, 134: 151–157. <https://doi.org/10.1016/j.fitote.2019.02.019>
- [32] PEÑA-MORÁN O., VILLARREAL M., ÁLVAREZ-BERBER L., MENESES-ACOSTA A., 和 RODRÍGUEZ-LÓPEZ V. 法加罗氏菌中天然存在的鬼臼毒素的细胞毒性、治疗后回收率和选择性分析。乳腺癌细胞系上的法加罗伊斯。分子, 2016, 21(8): 1013. <https://doi.org/10.3390/molecules21081013>
- [33] LÓPEZ-LÁZARO M. 我们应该筛选多少次化学文库才能发现一种抗癌药物? 今天的药物发现, 2015, 20(2): 167–169. <https://doi.org/10.1016/j.drudis.2014.12.006>
- [34] UDAYA SANKAR K. 提取过程。在: 传统和先进的食品加工技术。约翰威利父子公司, 奇切斯特, 2014: 129–158.
- [35] JUKIĆ I., POŽAR M., 和 LOVRINČEVIĆ B. 水溶液和非水溶液中乙醇动力学的比较分析。物理化学化学物理, 2020, 22(41): 23856–23868. <https://doi.org/10.1039/D0CP03160G>
- [36] DIBACTO R. E. K., TCHUENTE B. R. T., NGUEDJO M. W., TIENCHEU Y. M. T., NYOBE E. C., EDOUN F. L. E., KAMINI M. F. G., DIBANDA R. F., 和 MEDOUA G. N. 喀麦隆食用的部分美洲珍珠皮的总多酚和黄酮含量及抗氧化能力。科学世界杂志, 2021, 2021: 1–11. <https://doi.org/10.1155/2021/8882594>
- [37] KAYAHAN S., 和 SALOGLU D. 生和熟土耳其朝鲜蓟品种的酚类化合物和抗氧化活性的比较。可持续粮食系统前沿, 2021, 5. <https://doi.org/10.3389/fsufs.2021.761145>
- [38] SYABANA M. A., YULIANA N. D., BATUBARA I., 和 FARDIAZ D. 印度尼西亚常用蔬菜和香料的抗糖尿病活性筛选和核磁共振谱。食品科技, 2021, 41(1): 254–264. <https://doi.org/10.1590/fst.14120>
- [39] INDRAYANTO G., PUTRA G. S., 和 SUHUD F. 体外生物测定方法的验证: 在草药研究中的应用。原料药、辅料和相关方法学概况, 2021, 46: 273–307. <https://doi.org/10.1016/bs.podrm.2020.07.005>
- [40] PRITCHETT J. C., NAESENS L., 和 MONTOYA J. 治疗人类疱疹病毒-6感染。在: 人类疱疹病毒人类疱疹病毒-6A、人类疱疹病毒-6B和人类疱疹病毒-7。爱思唯尔, 阿姆斯特丹, 2014: 311–331.
- [41] WEERAPREEYAKUL N., NONPUNYA A., BARUSRUX S., THITIMETHAROCH T., 和 SRIPANIDKULCHAI B. 评估六种草药对肝癌细胞系的抗癌潜力。中医, 2012, 7(1): 15. <https://doi.org/10.1186/1749-8546-7-15>