

Open Access Article



<https://doi.org/10.55463/issn.1674-2974.49.12.15>

New Record of Marine Wood-Borer Species in Mangrove Forest of Nusa Lembongan, Bali, Indonesia

I. Ketut Ginantra^{1*}, I. Made Saka Wijaya¹, Ida Ayu Astarini^{1,2}, Khairul Adha A. Rahim³

¹ Biology Study Program, Faculty of Mathematics and Natural Sciences, Udayana University, Bali, Indonesia

² Master of Environmental Science Study Program, Postgraduate Program, Udayana University, Bali, Indonesia

³ Faculty of Resources Science and Technology, University Malaysia Sarawak, Malaysia

* Corresponding author: ketut_ginantra@unud.ac.id

Received: July 13, 2022 / Revised: September 15, 2022 / Accepted: November 11, 2022 / Published: December 30, 2022

Abstract: Research has been carried out on the presence of wood-borer species in the mangrove forest of Nusa Lembongan, in May-June 2022. The study was conducted at five representative sites of mangrove forests in Nusa Lembongan. The data collection of the mangrove wood-borer was carried out by purposive sampling, which directly searches for dead mangrove stems, twigs, or roots that were attacked by the marine wood-borer found at each site. A total of There were new record four species of mangrove wood-borer, namely, *Phaenops* sp. beetle group (Coleoptera) and *Cenoloba* sp. moth group (Lepidoptera), two species of Mollusks, namely, *Bankia minima* and *Bactronophorus* sp. were identified. The larval stages of *Phaenops* sp. were found within the dead mangrove stems of *Rhizophora apiculata* and *Sonneratia alba*. *Cenoloba* sp. which was found in the larvae phase, the host plant was the dead stems of the mangrove *Sonneratia alba*. *Bankia minima*, the host plants are dead stems of mangroves *Sonneratia alba*, *Rhizophora apiculata*, *Rhizophora stylosa* and *Avicennia* sp. *Bactronophorus* sp., the host plant is dead wood from the mangrove *Sonneratia alba*, *Rhizophora apiculata*. The most widely distributed wood-borer species in the mangroves of Nusa Lembongan is *Bankia minima*. The results showed that four species of marine wood-borer are new records in the mangrove forest of Nusa Lembongan.

Keywords: marine wood-borers, *Phaenops* sp., *Bankia minima*, mangrove.

印度尼西亚巴厘岛蓝梦岛红树林中海洋蛀木虫的新记录

摘要: 2022 年 5 月至 6 月, 对蓝梦岛红树林中蛀木虫物种的存在进行了研究。该研究是在蓝梦岛红树林的五个代表性地点进行的。红树林蛀木虫的数据收集是通过有目的的抽样进行的, 即直接搜索在每个地点发现的被海洋蛀木虫袭击的死红树茎、树枝或根。共有 4 种新记录的红树蛀木虫, 即褐藻。甲虫组(粉翅鸡)和芹菜。蛾类(鳞翅目), 两种软体动物, 即最小的银行和杆菌属。被确定。褐藻。的幼虫阶段。在根茎菌和白铃的死红树茎中发现。切诺洛巴服务提供商。在幼虫阶段被发现, 寄主植物是红树林海桑的枯死茎。最小的银行, 寄主植物是红树林海桑、细根根瘤菌、柱状根瘤菌和白骨。的枯茎。杆菌属, 寄主植物是来自红树林白铃, 根茎菌的枯木。蓝梦岛红树林中分布最广的蛀木虫物种是最小的银行。结果表明, 四种



海洋蛀木虫为蓝梦岛红树林新记录种.

关键词：海洋蛀木虫、酚类、最小的银行、红树林.

1. Introduction

The mangrove ecosystem is a system comprising of organisms (animals, plants, microbes) that interact with environmental factors in a mangrove habitat. The mangrove ecosystem is a unique ecotone (transitional area), which connects the life of land and sea biota [1]. The Nusa Lembongan mangrove forest with a forest area of 202 ha. some are planted and most of them are natural forests. The diversity of mangrove flora in Nusa Lembongan consists of three main components, namely, major, minor and associated mangroves. Approximately, around 13 true mangrove species and several associated plant species have been found in the Nusa Lembongan mangroves. True mangrove species include *Rhizophora* spp., *Avicennia* spp. *Sonneratia alba*, *Lumnitzera racemosa*, *Xylocarpus granatum* and associated species include *Pandanus tectorius*, *Hibiscus tiliaceus*, *Calotropis gigantea* and *Ipomoea pes-caprae* [2, 3].

Mangroves are the habitat of several aquatic fauna (mollusks, fish, crustaceans, insects) and avifauna (birds). Mangrove wood-borer is one of the fauna groups that use mangrove plants for their life. The marine wood-borer consists of various species, the fauna groups identified are from the Mollusks, Crustacean and Insect groups. Mollusks group, the best known are *Teredo*, *Bankia* and *Martesia*, Crustacean group includes *Limnoria*, *Chelura* and *Sphaeroma*; and Insect group (moth and beetle larvae), common genera are *Bottegia*, *Cenoloba*, and *Phaenops*.

The distribution area of the sea wood-borer is very wide and is found in almost all saltwater and brackish waters, in tropical waters, these animals can thrive and can be found throughout the year [4, 5]. Shipway et al. [6] and Weigelt et al. [3] stated that the marine wood-borer worm is a group of drill and wood eating bivalves that cause structural damage to mangrove trunks or roots. The hosts of several types of wood-borers that have been found are dead logs of the mangrove species *Rhizophora* spp., *Sonneratia* spp., and *Avicennia* spp.

Several studies on the marine wood-borer in mangrove forests of Southeast Asia have been conducted, such in Kalimantan, Peninsular Malaysia, Taiwan, China. Diba et al. [7] found several species of wood-borers including *Neoteredo reynes*, *Teredo pocalifer*, *Teredo utriculus*, *Teredo siamensis*, *Teredo navalis*, *Teredo batiliformis*, *Teredo dagmarae*, *Teredo dallii*, *Martesia setacea*, *Bankia setacea*, *Bankia minima* and *Petricola pholadiformis*; from the Teredinidae and Pholadidae families in the mangrove forests of West Kalimantan, Loo et al. [8] identified the

species *Dicyathifer mannii* and *Bankia gracilis* (Mollusca: Teredinidae) in the mangrove area of the Kuala Penyu area of Malaysia; Loo et al. [8] identified the species *Bactronophorus thoracites* and *Bankia gracilis* from Sabah Waters, Malaysia. Davidson et al. [9], Thiri and Yang [10] and Trinh et al. [11] reviewed the wood boring Isopod pest, *Sphaeroma* spp., which causes a destructive impact on Mangrove Forest in China, Southwestern Taiwan and Binh Dinh province, Vietnam; and Treneman et al. [12] found Six shipworm species in the coastal waters of Japan, including *Bankia* spp., *Teredothyra smithi*, *Psiloteredo* sp., *Lyrodus takanoshimensis*, and *Teredo* spp. and *Uperotus clava*. Some research results show that the presence of woodborer pests can cause damage to mangroves. Marine woodborers can negatively impact mangroves and change the morphology of the tree.

Damage to mangrove roots (pneumatophores and lenticels), namely with more necrotic tissue and holes in tree trunks, which resulted in the collapse of *Avicennia marina* mangrove plants. The hollow roots of *Rhizophora stylosa* cause the collapse of the mangrove tree [9].

Research data on the diversity of wood-borer species in mangroves in Indonesia is still very minimal. Indonesia, *Phaenops* sp, *Cenoloba* sp, *Bankia minima* and *Bactronophorus* sp. were reported previously this study, we provide the first report on the occurrence of marine wood-borer in Nusa Lembongan, Bali, Indonesia. In fact, the scientific data regarding wood-borer in the Nusa Lembongan mangroves, are not yet available. In the Mangrove Forest of Nusa Lembongan, twigs, stems and roots of dead mangroves places were found, as well as showing signs of the presence of wood-borers in them. Thus, this study aims to identify wood-borer species in several species of mangrove plants and the results of this study will be a new record for the existence of wood-borers in the mangroves of Nusa Lembongan.

2. Materials and Methods

2.1. Study Sites

The research was conducted for two months, from May-June 2022 in the mangrove ecosystem area of Nusa Lembongan. Table 1 shows the characteristics, GPS reading (coordinates) and the description of study sites in mangrove forest Nusa Lembongan and study sites are shown in Figure 2. Identification of mangrove wood-borer species was carried out in the Animal Taxonomy Laboratory, Biology, Faculty of

Table 1 The coordinates and description of study sites in mangrove forest Nusa Lembongan [13]

Location	Coordinates	Dominance species
Site 1	8°41'17.20" S 115°27'19.31" E	<i>Rhizophora apiculata</i> , <i>Rhizophora stylosa</i> , <i>Sonneratia alba</i>
Site 2	8°40'55.62" S 115°27'44.92" E	<i>Rhizophora stylosa</i> , <i>Bruguiera gymnorhiza</i> , <i>Sonneratia alba</i>
Site 3	8°40'16.12" S 115°27'54.45" E	<i>Bruguiera gymnorhiza</i> , <i>Ceriops tagal</i> , <i>Rhizophora stylosa</i> , <i>Xylocarpus granatum</i> , <i>Avicennia rumphiana</i> , <i>Rhizophora x lamarckii</i> , <i>Sonneratia alba</i>
Site 4	8°40'02.82" S 115°27'41.39" E	<i>Rhizophora stylosa</i> , <i>Ceriops tagal</i> , <i>Xylocarpus granatum</i> , <i>Avicennia marina</i> , <i>Sonneratia alba</i> .
Site 5	8°40'11.99" S 115°28'16.28" E	<i>Rhizophora apiculata</i> , <i>Rhizophora stylosa</i> , <i>Sonneratia alba</i>



Fig. 1 Maps (Google Earth) showing sampling stations at five locations in the mangrove forest at Nusa Lembongan

2.2. Data Collection and Species Identification

The data collection of the mangrove wood-borer was carried out by purposive sampling [14]. The sample collection was limited to dead mangrove stems at five research sites. The inventory was carried out by directly searching for dead mangrove stems, twigs, or roots that were attacked by the marine wood-borer found in each site (5 sites) of the Nusa Lembongan mangrove forest.

All affected wood is taken and collected. Wood that is attacked by the wood-borer is cut with an ax to a size of about 1 meter, then split using an ax to remove the marine wood-borer. All marine wood-borer specimens found in the wood were taken with tweezers and collected in vials 70% alcohol. Identification of marine wood-borer species is based on morphological characteristics and shell shape (for mollusks) and characteristics of holes in dead mangrove wood [7, 11, 15-18].

2.3. Data Analysis

The data were analyzed descriptively, by describing the morphological structures, shell structure, holes in the dead mangrove wood. The specimens found were compared with the descriptions of the species or genus [7, 8, 10, 16-19].

3. Results and Discussion

3.1. Existence of Wood-Borer in Mangrove Forest

A total of four species of marine wood-borer in Nusa Lembongan were recorded. This includes two species from the insect class, *Phaenops* sp. (Coleoptera, beetle) and *Cenoloba* sp. (Lepidoptera, moths), respectively. Both species are found in the larval phase in *R. stylosa* and *S. alba*, which were spread out at sites 1, 3 and 4. The other two marine wood-borer species are of the Bivalvia, namely, *Bankia minima* and *Bactronophorus* sp. These two species were found mainly in *Sonneratia alba*, *Rhizophora apiculata* and *Avicennia* sp and *Bankia minima* is the most widely distributed species in the Nusa Lembongan mangroves (Table 2).

Previous studies on species composition and distribution of fauna in mangrove forests in Indonesia mostly focused on crabs, shellfish (Bivalvia), snails (Gastropod), fish and birds [13-18, 21-23]. However, the information on the occurrences and species composition of marine wood-borer in mangrove areas is still lacking. Thus, the discovery of four species of marine wood-borer mangrove areas is the first record of these species on the island of Nusa Lembongan, Bali, Indonesia. The larval characteristics of *Phaenops* sp. are similar to those of the same genus described by Ciesla [17]. The larvae phase of this genus generally attacks dead wood cores from mangrove plants, including *Rhizophora* sp., *Sonneratia* sp. The adult phase of this beetle lays its eggs in the pith of the mangrove wood, then hatches into larvae that can make holes in the wood. The larval characteristics of *Cenoloba* sp. found in the mangroves of Nusa

Lembongan correspond/resemble with to the characteristics of the same genus described by Abu El-Ghiet [16]. It was further stated that the species *Cenoloba oblitalis* is a pest that attacks mangrove plants (references). This species, in the larval phase,

mostly attacks the *Avicennia marina* mangroves, especially on the fruit and leaves. Abdel-Rahman et al. [24] also found other insect species in the larval stage, namely, *Trichogramma* sp., which is a common parasitoid in *Avicennia marina* mangroves.

Table 2 Marine wood-borer species were collected from five sites in Nusa Lembongan mangrove forest

No.	Species (common name)	Host plant	Class/Order/ Family	Occurrence on site				
				1	2	3	4	5
1	<i>Phaenops</i> sp. (Larvae phase of Flatheaded wood-borer)	<i>Rhizophora stylosa</i> , <i>Sonneratia alba</i>	Insecta/Coleoptera/Buprestidae			V	V	
2	<i>Cenoloba</i> sp. (Larvae phase of moth)	<i>Sonneratia alba</i>	Insecta/Lepidoptera/Tineodidae	V		V		
3	<i>Bankia minima</i> (Shipworm wood-borer)	<i>Sonneratia alba</i> , <i>Rhizophora stylosa</i> , <i>Rhizophora apiculata</i> , <i>Avicennia</i> sp.	Bivalvia/Myoida/Teredinidae	V	V		V	V
4	<i>Bactronophorus</i> sp. (woodworm wood-borer "Tembelo")	<i>Sonneratia alba</i> , <i>Rhizophora apiculata</i>	Bivalvia/Myoida/Teredinidae			V	V	

Wood-borer from the Lepidoptera and Coleoptera groups caused damage to the mangrove wood during the larval stage. This attack begins when the female insect lays its eggs into the bark. Eggs are laid or oviposition in several different parts of the stem (2–6 clusters), each cluster can contain 100–200 eggs. After the eggs hatch into larvae, they begin eating wood while forming a tunnel. Damage to the wood occurs after being drilled by the larvae, which is indicated by discoloration and the stem begins to weaken. The tunnels formed by these larvae cause the tree to weaken structurally, which can then become brittle [25].

Bankia minima is wood-borer worm from the mollusk group, which is common species found in the mangroves of Nusa Lembongan. The host of this mollusk is dead mangrove wood, namely from the species *Sonneratia alba*, *Rhizophora apiculata*, *Rhizophora stylosa* and *Avicennia marina*. Several previous researchers have also published the existence of *Bankia* sp. on mangrove wood in some mangrove forest areas. Diba et al. [7] noted the presence of *B. setacea* and *B. minima* in the mangrove forests of West Kalimantan. Treneman et al. [12] reported the presence of *Bankia* spp. in the coastal mangrove forests of Japan. Loo et al. [8] and Loo et al. [19] found *Bankia gracilis* in the mangrove forest Kuala Penyu Malaysia. *Bactronophorus* sp. is also a wood-borer from the mollusk group, which is less common in the mangroves of Nusa Lembongan.

Species of the same genus, *Bactronophorus thoracites*, have been reported by Loo et al. [8] and Loo et al. [20] in the Kinabalu mangrove forest, Sabah, Malaysia. Hendy and Cragg [6] have also studied the impact of the mollusk group (Family Teredinidae) wood-borer on damage to mangrove wood in the *Rhizophora stylosa* species.

The ability of wood-borers to digest wood or drill wood, which generally contains cellulose, is greatly assisted by the symbiosis with microorganisms. Some species of wood-borer in the digestive tract there are bacteria and protozoa. Most wood-borers (groups of lepidoptera, coleoptera, molluscs, isopods) can digest wood because of their symbiosis with bacteria and protozoa in their digestive tract. Only a few species of wood-borers can actually produce their own cellulose enzymes. In several shipworm (Trinididae) species, dense communities of intracellular bacterial endosymbionts have been observed within specialized cells (microorganisms) of the gills. These bacteria are proposed to contribute to the digestion of wood by the host [26, 27]. Altamia et al. [28] have also isolated three bacterial species, namely, *Teredinibacter turnerae*, *Teredinibacter purpureus* and *Teredinibacter franksiae*, in the digestive tract of shipworm species *Bankia setacea* (Bivalvia: Teredinidae). It was further stated that these bacteria become endosymbiotic in digestion and play a role in producing cellulose enzymes to digest wood cellulose.

3.2. Description of Wood-Borer Species

3.2.1. *Phaenops* Sp. (Coleoptera, Buprestidae)

Larvae phase: The body of the larvae is white, legless, similar to the larvae of the bark beetle, but the shape of the body and the head are slightly different from the larvae of the bark beetle, namely the body shape is longer, the head (cephal) and the body segment of the thorax are much wider than the next segment, also usually has a hardened plate at the top and bottom (Figure 2).

The larvae immediately drill into the wood and slowly enter the core, making it difficult to see. The

larvae will continue to consume wood, which is the brand's only food source. Inside the wood, the larvae become safe, continue to dig for food for several years. Adult: the adult phase is a flat-headed wood-borer, a distinctive oval body shape, small antennae. Its body size is relatively small to relatively large, which is between 6–64 mm. Adult Buprestidae are called metallic wood-borers because they are iridescent or metallic looking underneath and sometimes on top. Adult beetles will lay their eggs in cracks in wooden objects, floorboards and timbers [29-31].



Fig. 2 *Phaenops* sp. (larvae phase); image on the left: larval morphology; on the right: larvae in the wood core hole [15]

3.2.2. *Cenoloba* Sp. (Lepidoptera, Tineodidae)

Larvae: Larvae eruciform, peripneustic, frequently with 8 pairs of limbs. All larvae of moths and butterflies (Lepidoptera) have proles on their abdomen. These legs are tipped with hooks, the crochet (Figure 3). Pupae usually adecticous and more or less object, and generally enclosed in a cocoon or an earthen cell; a few primitive forms are decticous and exarate. Adult: Insects with 2 pairs of membranous wings; cross-veins few. The body, wings and appendages were clothed with broad scales. Mandibles are almost always vestigial or absent, and the principal mouth parts generally represented by a suctorial proboscis formed by the maxillae. Wing tracheation complete [16].



Fig. 3 *Cenoloba* sp. (larvae phase) [15]

3.2.3. *Bankia Minima* (Mollusca, Teredinidae)

The marine wood-borer found had a length of about 4–7 cm, its body color was brownish, somewhat similar to the color of its host plant wood, its tail color was yellowish white to brown and had a soft and transparent body. Sea wood-borer worms contain water, but the water is easily reduced when the wood-borer's body is removed from the wood. The shape will immediately change to small even though when first encountered it is countless. The head has a pair of small spheres called shells, white to brown in color, which are useful for making holes in wood. The tail shape (palette) is used for species identification. Inside

the mollusk's wood it forms a circular tunnel in cross-section and is lined with a calcareous material extruded by the mollusk (Figure 4).




Fig. 4 *Bankia* sp. (left image: body morphology; right image: circular tunnel in dead wood) [15]

3.2.4. *Bactronophorus* Sp. (Mollusca, Teredinidae)

The body of *Bactronophorus* sp. tubular, its body length reaches 4 cm, pale white. The body can be elongated or shortened. The wider anterior end is closed, has a rounded end. From there, the tube tapers to the open posterior end, by the central septum. The siphon projects through this end to feed and breathe. They can be pulled into the tube and ends can be sealed with a set of pallets. The mollusk's two tiny valves are inside the tube along with the mantle, intestines, and other soft organs. Inside this mollusk wood also forms a circular tunnel in its cross-section and is coated with calcareous material produced by the mollusk (Figure 5). The larger species of sea woodwork, which also lives in dead mangrove trunks, is often called “Tembelo” by the people of Eastern Indonesia and can be consumed.



Fig. 5 *Bactronophorus* sp. (left image: body morphology; right image: Tunnel in log) [15]

4. Conclusion

Four species of mangrove wood-borer were found in the Nusa Lembongan Mangrove Forest. Two species were found in the larvae phase, namely *Phaenops* sp. from the beetle (Coleoptera) group, the host plant is dead mangrove stems from *Rhizophora apiculata*, *Sonneratia alba* and *Cenoloba* sp. is a group of moths (Lepidoptera), whose host plant is the dead stems of the mangrove *Sonneratia alba*. Two species from the mollusk group, namely *Bankia minima*, whose host plant is dead stem from the mangrove *Sonneratia alba*, *Rhizophora apiculata*, *Rhizophora stylosa*, *Avicennia* sp. and *Bactronophorus* sp., the host plant is dead wood from the mangrove *Sonneratia alba*, *Rhizophora apiculata*. *Bankia minima* is the most widely distributed species among the mangroves of Nusa Lembongan. Research on the existence of marine wood-borers in mangrove ecosystems in Bali has never been reported before, so the discovery of wood-borers,

namely, two species from the mollusk group and two species from the insect group, is a recent finding.

Acknowledgments

We would like to thank the Research and Community Service Institute of Udayana University, for funding research through the Foreign Cooperation Research Grant scheme. Thanks to Abdi Janaguna, Sugiyama, Adianto Wibisana and Made Goura Premananda for their assistance in field sampling.

References

[1] KITAMURA S., ANWAR C., CHANIAGO A. and BABA S. *Handbook of Mangroves in Indonesia - Bali & Lombok*. International Society for Mangrove Ecosystems, Tokyo, 1997.
http://www.mangrove.or.jp/img/publications/book_pdf/C-PB01-Handbook.pdf

[2] PALGUNA I. B. A., ARDHANA I. P. G. and ARTHANA I. W. Mangrove Forest Structure and Diversity In Nusa Lembongan, Nusa Penida Sub District, Klungkung District. *Ecotrophic*, 2017, 11(2): 109-115.
<https://doi.org/10.24843/EJES.2017.v11.i02.p07>

[3] PRICILLIA C. C., PATRIA M. P. and HERDIANSYAH H. Environmental conditions to support blue carbon storage in mangrove forest: A case study in the mangrove forest, Nusa Lembongan, Bali, Indonesia. *Biodiversitas*, 2021, 22(6): 3304-3314. <https://doi.org/10.13057/biodiv/d220636>

[4] JENOH E. M., ROBERT E. M. R., LEHMANN I., KIOKO E., BOSIRE J. O., NGISIANGE N., DAHDUHGUEBAS and KOEDAM N. Wide Ranging Insect Infestation of the Pioneer Mangrove *Sonneratia alba* by Two Insect Species along the Kenyan Coast. *PLoS ONE*, 2016, 11(5): e0154849.
<https://doi.org/10.1371/journal.pone.0154849>

[5] PALVAST P. & VELDE G. New threats of an old enemy: The distribution of the shipworm *Teredo navalis* L. (*Bivalvia*: *Teredinidae*) related to climate change in the Port of Rotterdam area the Netherlands. *Marine Pollution Bulletin*, 2011, 62(8): 1822-1829.
<https://doi.org/10.1016/j.marpolbul.2011.05.009>

[6] HENDY I. W. & CRAGG S. M. Rhizophora stylosa prop roots even when damaged prevent wood-boring teredinids from toppling the trees. *Hydrobiologia*, 2017, 803: 333-344.
<https://doi.org/10.1007/s10750-017-3106-6>

[7] DIBA F., WANAMUKTI B., ADHA K. and CHENG-ANN C. Species Diversity of Marine Wood Borer Worms from the Setapok Besar Mangrove Forest, Singkawang City, West Kalimantan. *Jurnal Enggano*, 2021, 6(1): 62-79.
<https://doi.org/10.31186/jenggano.6.1.62-79>

[8] LOO Z. A., CHENG-ANN C., RAHIM K. A. A. and DIBA F. First record of marine wood borer (Mollusca: *Teredinidae*) *Dicyathifer mannii* Wright (1866) in Sabah, Malaysia, with detailed measurement metrics. *Borneo Journal of Marine Science and Aquaculture*, 2019, 03(1): 37- 40. <https://doi.org/10.51200/bjomsa.v3i1.1889>

[9] DAVIDSON T. M., DE RIVERA C. E. and HWEY-LIAN H. Damage and alteration of mangroves inhabited by a marine wood-borer. *Marine Ecology Progress Series*, 2014, 5(16): 177-185. <https://doi.org/10.3354/meps11046>

[10] THIRI M. & YANG Y. Review on Possible Factors for

Outbreak of Wood Boring Isopod, *Sphaeroma* spp. Which Causes Destructive Impact on Mangrove Forest in China. *Open Journal of Ecology*, 2022, 12(3): 211-235.
<https://doi.org/10.4236/oje.2022.123013>

[11] TRINH V. H., NGUYEN Q. H., TUAT L. V., TRA T. L. and DANG N. B. New record of a wood-boring isopod damaged *Sonneratia alba* J. Sm. in Thi Nai lagoon, Binh Dinh province, Vietnam. *International Journal of Technical & Scientific Research Engineering*, 2019, 2(3): 12-18.
<https://ijtsre.org/papers/2019/ev2c3/68745214.pdf>

[12] TRENEMAN N. C., CARLTON J. T., BORGES L. M. S., SHIPWAY J. R., RAUPACH M. J. and ALTERMARK B. Species diversity and abundance of shipworms (Mollusca: Bivalvia: *Teredinidae*) in woody marine debris generated by the Great East Japan Earthquake and Tsunami of 2011. *Aquatic Invasions*, 2018, 13(1): 87-100.
<https://doi.org/10.3391/ai.2018.13.1.07>

[13] GINANTRA I. K., MUKSIN I. K. and JONI M. Crab diversity as support for ecotourism activities in Pejarakan Mangrove Forest, Buleleng, Bali, Indonesia. *Biodiversitas*, 2021, 22(10): 4139-4145.
<https://doi.org/10.13057/biodiv/d221003>

[14] ETIKAN I., MUSA S. A. and ALKASSIM R. S. Comparison of Convenience Sampling and Purposive Sampling. *American Journal of Theoretical and Applied Statistics*, 2015, 5(1): 1-4.
<https://doi.org/10.11648/j.ajtas.20160501.11>

[15] GINANTRA I. K., MUKSIN I. K., SUASKARA I. D. M. and JONI M. Diversity and distribution of mollusks at three zones of mangrove in Pejarakan, Bali, Indonesia. *Biodiversitas*, 2020, 21(10): 4636-464.
<https://doi.org/10.13057/biodiv/d211023>

[16] ABU EL-GHIET U. M. A New Record for Mangrove Plume Moth *Cenoloba oblitalis* (Walker, 1864) (Fam. *Tineodidae*: *Lepidoptera*) Attacking Mangrove Plant in Saudi Arabia. *Journal of Plant Protection and Pathology, Mansoura University*, 2019, 10(2): 97-100.
<https://doi.org/10.21608/jppp.2019.40883>

[17] CIESLA W. M. *Woodborer, Metallic Woodboring Beetle*. Forest Health Management International, 2018.
<https://texasinsects.tamu.edu/woodborer-metallic-woodboring-beetle/>

[18] VELÁSQUEZ M., SHIPWAY J. R., LIRA C., CAPELO J. and NARCISO S. Shipworms from Venezuela (Mollusca, Bivalvia, *Teredinidae*): an updated survey. *The Festivus*, 2017, 49(4): 302-316.
<https://doi.org/10.54173/F494302>

[19] SARI A., TUWO A., SARU A. and RANI C. Diversity of fauna species in the mangrove ecosystem of Youtefa Bay Tourism Park, Papua, Indonesia. *Biodiversitas*, 2022, 23(9): 4490-4500. <https://doi.org/10.13057/biodiv/d230915>

[20] LOO Z. A., CHENG-ANN C., RAHIM K. A. A. and DIBA F. Redescription of *Bactronophorus thoracites* Gould (1856) and *Bankia gracilis* Moll (1935) from Sabah Waters, Malaysia, with Short Ecological Notes and Measurements Metrics. *Scientific Journal of Fisheries and Marine*, 2021, 14(1): 150-159. <https://doi.org/10.20473/jipk.v14i1.26825>

[21] ATLANTA V., AMBARWATI R., RAHAYU D. A. and MUJIONO N. Diversity of bivalves on the North Coast of Lamongan, East Java, Indonesia. *Biodiversitas*, 2022, 23(8): 4263-4271. <https://doi.org/10.13057/biodiv/d230850>

[22] BADERAN D. W. K., HAMIDUN M. S., UTINA R., RAHIM S. and DALI R. The abundance and diversity of

Mollusks in mangrove ecosystem at coastal area of North Sulawesi, Indonesia. *Biodiversitas*, 2019, 20(4): 987-993. <https://doi.org/10.13057/biodiv/d200408>

[23] DEWIYANTI I. & KARINA S. Diversity of Gastropods and Bivalves in mangrove ecosystem rehabilitation areas in Aceh Besar and Banda Aceh districts, Indonesia. *Aquaculture, Aquarium, Conservation & Legislation. International Journal of the Bioflux Society*, 2012, 5(2): 55-59. http://www.bioflux.com.ro/docs/AACL_5.2.1.pdf

[24] ABDEL-RAHMAN R. G., AGAMY E., MOHAMMED F., USAMA M. and EL GHIET A. Widespread of *Trichogramma* egg parasitoids among mangrove forests, *Avicennia marina*. *Egyptian Journal of Desert Research*, 2019, 69: 165-173. <https://doi.org/10.21608/ejdr.2021.18001.1061>

[25] HANNON E. & BROWN J. J. *Carpenterworm Moth Insect Pest Management in Hybrid Poplars Series*. Washington State University, Washington DC, 2017. <https://doi.org/10.13140/RG.2.2.10827.49447>

[26] BETCHER M. A., FUNG J. M., HAN A. W., O'CONNOR R., SERONAY R., CONCEPCION G. P., DISTEL D. L. and HAYGOOD M. G. Microbial Distribution and Abundance in the Digestive System of Five Shipworm Species (Bivalvia: Teredinidae). *PLoS ONE*, 2012, 7(9): e45309. <https://doi.org/10.1371/journal.pone.0045309>

[27] SHIPWAY J. R., ALTAMIA M. A., ROSENBERG G., CONCEPCION G. P. and HAYGOOD M. G. A rock-boring and rock-ingesting freshwater bivalve (shipworm) from the Philippines. *Proceedings of the Royal Society*, 2019, 286(1905): 1-10. <https://doi.org/10.1098/rspb.2019.0434>

[28] ALTAMIA M. A., SHIPWAY J. R., STEIN D., BETCHER M. A., FUNG J. M., JOSPIN G., EISEN J., HAYGOOD M. G. and DISTEL D. L. *Teredinibacter haidensis* sp. nov., *Teredinibacter purpureus* sp. nov. And *Teredinibacter franksiae* sp. nov., marine, cellulolytic endosymbiotic bacteria isolated from the gills of the woodboring mollusc *Bankia setacea* (Bivalvia: Teredinidae) and emended description of the genus *Teredinibacter*. *International Journal of Systematic and Evolutionary Microbiology*, 2021, 71(004627): 1-11. <https://doi.org/10.1099/ijsem.0.004627>

[29] PURYONO S. & SURYANTI S. Gastropod Diversity in Mangrove Forests of Mojo Village, Ulujami District, Pemalang Regency, Indonesia. *Journal of Ecological Engineering*, 2019, 20(1): 165-173. <https://doi.org/10.1xxix111/2xxix198993/93940>

[30] TAN L. W. H. & PETER K. L. N. 2019. *A Guide to Seashore Life: Wood Borer*. Singapore Science Centre, Singapore, 2019. <http://mangrove.nus.edu.sg/pub/seashore/text/261.htm>

[31] WEIGELT R., LIPPERT H., BORGES L. M. S. and BASTROP R. First time DNA barcoding of the common shipworm *Teredo navalis* Linnaeus 1758 (Mollusca: Bivalvia: Teredinidae): Molecular-taxonomic investigation and identification of a widespread wood-borer. *Journal of Experimental Marine Biology and Ecology*, 2016, 475: 154-162. <https://doi.org/10.1016/j.jembe.2015.11.008>

参考文献:

[1] KITAMURA S., ANWAR C., CHANIAGO A. 和 BABA S. 印度尼西亚红树林手册 - 巴厘岛和龙目岛。国

际红树林生态系统协会，东京，1997. http://www.mangrove.or.jp/img/publications/book_pdf/C-PB01-Handbook.pdf

[2] PALGUNA I. B. A., ARDHANA I. P. G., 和 ARTHANA I. W. 蓝梦岛、珀尼达岛分区、克隆孔区的红树林结构和多样性。生态营养的, 2017, 11(2): 109-115. <https://doi.org/10.24843/EJES.2017.v11.i02.p07>

[3] PRICILLIA C. C., PATRIA M. P. 和 HERDIANSYAH H. 支持红树林蓝碳储存的环境条件：以印度尼西亚巴厘岛蓝梦岛红树林为例。生物多样性, 2021, 22(6): 3304-3314. <https://doi.org/10.13057/biodiv/d220636>

[4] JENOH E. M., ROBERT E. M. R., LEHMANN I., KIOKO E., BOSIRE J. O., NGISIANGE N., DAHDOUNH-GUEBAS 和 KOEDAM N. 肯尼亚海岸沿线两种昆虫对先驱红树林白海桑的大范围昆虫侵扰。一号的公共科学图书馆一号, 2016, 11(5): e0154849. <https://doi.org/10.1371/journal.pone.0154849>

[5] PALVAST P. 和 VELDE G. 老敌人的新威胁：与荷兰鹿特丹港地区气候变化有关的船虫海军打磨。(双壳纲：蟹科)的分布。海洋污染公报, 2011, 62(8): 1822-1829. <https://doi.org/10.1016/j.marpolbul.2011.05.009>

[6] HENDY I. W. 和 CRAGG S. M. 柱状根茎菌的支柱根即使在损坏时也能防止钻木的土鳖类倒塌树木。水生生物学, 2017, 803: 333-344. <https://doi.org/10.1007/s10750-017-3106-6>

[7] DIBA F., WANAMUKTI B., ADHA K. 和 CHENG-ANN C. 西加里曼丹山口洋市文良港勿刹红树林海洋蛀虫的物种多样性。英语杂志, 2021, 6(1): 62-79. <https://doi.org/10.31186/jenggano.6.1.62-79>

[8] LOO Z. A., CHENG-ANN C., RAHIM K. A. A. 和 DIBA F. 马来西亚沙巴的海洋蛀木虫(软体动物：蟹科)曼尼双胞胎蠕(1866年)的首次记录，具有详细的测量指标。婆罗洲海洋科学与水产养殖杂志, 2019, 03(1): 37-40. <https://doi.org/10.51200/bjomsa.v3i1.1889>

[9] DAVIDSON T. M., DE RIVERA C. E. 和 HWEY-LIAN H. 海洋蛀木虫栖息的地方红树林的破坏和变化。海洋生态学进展丛书, 2014, 5(16): 177-185. <https://doi.org/10.3354/meps11046>

[10] THIRI M. 和 YANG Y. 蛀木等足类动物球形藻属爆发的可能因素综述 对中国红树林造成破坏性影响。开放生态杂志, 2022, 12(3): 211-235. <https://doi.org/10.4236/oje.2022.123013>

[11] TRINH V. H., NGUYEN Q. H., TUAT L. V., TRA T. L. 和 DANG N. B. 一种钻木等足类动物的新记录损坏了白海桑。在越南平定省的氏奈泻湖。国际技术与科学研究工程杂志, 2019, 2(3): 12-18. <https://ijtsre.org/papers/2019/ev2c3/68745214.pdf>

[12] TRENNEMAN N. C., CARLTON J. T., BORGES L. M. S., SHIPWAY J. R., RAUPACH M. J. 和 ALTERMARK B. 2011年东日本大地震和海啸产生的木质海洋垃圾中船虫(软体动物：双壳类：泥鳅科)的物种多样性和丰度。水生入侵, 2018, 13(1): 87-100. <https://doi.org/10.3391/ai.2018.13.1.07>

[13] GINANTRA I. K., MUKSIN I. K. 和 JONI M. 螃蟹多样性作为对印度尼西亚巴厘岛布勒冷佩拉坎红树林生态旅游活动的支持。生物多样性, 2021, 22(10): 4139-4145. <https://doi.org/10.13057/biodiv/d221003>

- [14] ETIKAN I., MUSA S. A. 和 ALKASSIM R. S. 便利抽样和目的抽样的比较。美国理论与应用统计杂志, 2015, 5(1): 1-4. <https://doi.org/10.11648/j.ajtas.20160501.11>
- [15] GINANTRA I. K., MUKSIN I. K., SUASKARA I. D. M. 和 JONI M. 印度尼西亚巴厘岛佩扎拉坎红树林三个区域软体动物的多样性和分布。生物多样性, 2020, 21(10): 4636-464. <https://doi.org/10.13057/biodiv/d211023>
- [16] ABU EL-GHIET U. M. 红树羽蛾金缕梅 (沃克, 1864 年) (翅科: 鳞翅目) 攻击沙特阿拉伯红树林植物的新记录。曼苏拉大学植物保护与病理学杂志, 2019, 10(2): 97-100. <https://doi.org/10.21608/jppp.2019.40883>
- [17] CIESLA W. M. 蛀木虫, 金属蛀木甲虫。森林健康管理国际, 2018. <https://texasinsects.tamu.edu/woodborer-metallic-woodboring-beetle/>
- [18] VELÁSQUEZ M., SHIPWAY J. R., LIRA C., CAPELO J. 和 NARCISO S. 来自委内瑞拉的船虫 (软体动物、双壳类、泥鳅科): 最新调查。节日, 2017, 49(4): 302-316. <https://doi.org/10.54173/F494302>
- [19] SARI A., TUWO A., SARU A. 和 RANI C. 在印度尼西亚巴布亚尤特法湾旅游公园红树林生态系统中, 动物物种是多样性。生物多样性, 2022, 23(9): 4490-4500. <https://doi.org/10.13057/biodiv/d230915>
- [20] LOO Z. A., CHENG-ANN C., RAHIM K. A. A. 和 DIBA F. 重新描述来自马来西亚沙巴水域的胸细菌(1856) 和银杏(1935), 附有简短的生态注释和测量指标。渔业和海洋科学杂志, 2021, 14(1): 150-159. <https://doi.org/10.20473/jipk.v14i1.26825>
- [21] ATLANTA V., AMBARWATI R., RAHAYU D. A. 和 MUJIONO N. 印度尼西亚东爪哇拉蒙甘北海岸双壳类动物的多样性。生物多样性, 2022, 23(8): 4263-4271. <https://doi.org/10.13057/biodiv/d230850>
- [22] BADERAN D. W. K., HAMIDUN M. S., UTINA R., RAHIM S. 和 DALI R. 印度尼西亚北苏拉威西沿海地区红树林生态系统中软体动物的丰度和多样性。生物多样性, 2019, 20(4): 987-993. <https://doi.org/10.13057/biodiv/d200408>
- [23] DEWIYANTI I. 和 KARINA S. 印度尼西亚亚齐大区和班达亚齐地区红树林生态系统恢复区腹足类和双壳类的多样性。水产养殖、水族馆、保护与立法。国际生物通量学会杂志, 2012, 5(2): 55-59. http://www.bioflux.com.ro/docs/AACL_5.2.1.pdf
- [24] ABDEL-RAHMAN R. G., AGAMY E., MOHAMMED F., USAMA M. 和 EL GHIET A. 赤眼蜂卵寄生蜂在红树林中广泛分布, 白骨码头。埃及沙漠研究杂志, 2019, 69: 165-173. <https://doi.org/10.21608/ejdr.2021.18001.1061>
- [25] HANNON E. 和 BROWN J. J. 杂交杨树系列木蛾害虫防治。华盛顿州立大学, 华盛顿特区, 2017. <https://doi.org/10.13140/RG.2.2.10827.49447>
- [26] BETCHER M. A., FUNG J. M., HAN A. W., O'CONNOR R., SERONAY R., CONCEPCION G. P., DISTEL D. L. 和 HAYGOOD M. G. 五种船虫 (双壳类: 船虫科) 消化系统中的微生物分布和丰度。第一科学公共图书馆, 2012, 7(9): e45309. <https://doi.org/10.1371/journal.pone.0045309>
- [27] SHIPWAY J. R., ALTAMIA M. A., ROSENBERG G., CONCEPCION G. P. 和 HAYGOOD M. G. 来自菲律宾的钻岩, 取食淡水双壳类(船水)。英国皇家学会会刊, 2019, 286(1905): 1-10. <https://doi.org/10.1098/rspb.2019.0434>
- [28] ALTAMIA M. A., SHIPWAY J. R., STEIN D., BETCHER M. A., FUNG J. M., JOSPIN G., EISEN J., HAYGOOD M. G. 和 DISTEL D. L. 海地珊瑚虫。十一月, 紫球菌。十一月 和 法兰西菌。11 月, 从钻木软体动物山竹 (双壳类: 蟹科) 的壳素分解内共生细菌, 并修订了泥杆菌属的描述。国际系统和进化微生物学杂志, 2021, 71(004627): 1-11. <https://doi.org/10.1099/ijsem.0.004627>
- [29] PURYONO S. 和 SURYANTI S. 印度尼西亚八马朗摄政区乌鲁贾米区莫乔村红树林中的腹足类动物多样性。生态工程学报, 2019, 20(1): 165-173. <https://doi.org/10.12911/22998993/93940>
- [30] TAN L. W. H. 和 PETER K. L. N. 2019. 海滨生活指南: 蛀木虫。新加坡科学中心, 新加坡, 2019. <http://mangrove.nus.edu.sg/pub/seashore/text/261.htm>
- [31] WEIGELT R., LIPPERT H., BORGES L. M. S. 和 BASTROP R. 常见船虫海蠊 1758 (软体动物: 双壳类: 蟹科) 的首次脱氧核糖核酸条形码: 分子分类学调查和广泛分布的蛀木虫的鉴定。实验海洋生物学与生态学杂志, 2016, 475: 154-162. <https://doi.org/10.1016/j.jembe.2015.11.008>